

Oral Health Fact Sheet for Medical Professionals

Children with Traumatic Brain Injury

Traumatic brain injury (TBI), a form of acquired brain injury, occurs when a sudden trauma causes damage to the brain. TBI can result when the head suddenly and violently hits an object, or when an object pierces the skull and enters brain tissue. Symptoms of a TBI can be mild, moderate, or severe, depending on the extent of the damage to the brain. (National Institute of Neurological Disorders and Stroke/NIH)

Oral Manifestations and Considerations

Oral

- Oral/dental trauma from TBI or self-injurious behaviors
- Bruxism
- GERD
- Inadequate oral hygiene due to cognitive impairments, spasticity and ataxia

Other Potential Disorders/Concerns

- Seizures
- Depression/Anxiety
- Post-traumatic stress disorder
- Personality disorders
- Substance abuse

Oral Side Effects of Commonly Prescribed Medications

Medication: Manifestations from TBI vary; therefore a range of medications may be prescribed:

SYMPTOM	MEDICATION	SIDE EFFECTS
	<i>Muscle Relaxants</i>	Xerostomia
Repetitive Behaviors	<i>Antidepressants</i>	Xerostomia, dysgeusia, stomatitis, gingivitis, glossitis, sialadenitis, bruxism, dysphagia, discolored tongue, oral edema
Aggressive Behaviors	A. <i>Anticonvulsants</i> (Dilantin)	A. Gingival hyperplasia, xerostomia, stomatitis, glossitis, dysgeusia
	B. <i>Antipsychotics</i>	B. Xerostomia, sialorrhea, dysphagia, dysgeusia, stomatitis, gingivitis, tongue edema, glossitis, discolored tongue
Hyperactivity	A. <i>Antihypertensive</i>	A. Xerostomia, dysphagia, sialadenitis, dysgeusia
	B. <i>CNS Stimulant</i>	B. Xerostomia

Children with Traumatic Brain Injury continued

Parent/Caregiver Support and Guidance

- Discourage consumption of cariogenic foods and beverages.
- Prescribe sugar-free medications if available.
- Recommend preventive measures such as topical fluoride and sealants.
- Advise the use of fluoridated toothpaste twice daily and support the family in following dental care instructions.
- Instruct caregiver on appropriate protocol following dental trauma (seek immediate professional care and locate/preserve missing tooth).
- Review safety issues appropriate to the age of the child, such as car seats, stair gates, bike helmets, and mouth guards to prevent oral-facial trauma.
- Discuss habits that may harm the child's teeth such as propping baby bottles, putting child to bed with bottle
- Refer to dentist any oral developmental abnormalities.

Further information (Medical Anticipatory Guidance, Dental Anticipatory Guidance, Oral Health Guidance for parents/Caregivers and Dental Professional Fact Sheet for Traumatic Brain Injury) can be found at:

http://dental.washington.edu/departments/omed/decod/special_needs_facts.php

References

- Taylor, HG., Yeates, KO., Wade, SL., Drotar, D., Stancin, T., Minich, N. (2002) A prospective study of short- and long-term outcomes after traumatic brain injury in children: behaviour and achievement. *Neuropsychology*, 16: 15–27.
- McKinlay, A., Grace, R.C., Horwood, L.J., Fergusson, D.M., Ridder, E.M., MacFarlane, M. (2008) Prevalence of traumatic brain injury among children, adolescents and young adults: Prospective evidence from a birth cohort. *Brain Injury*, 22(2): 175-81.

Additional Resources

- [NIH Institute for Traumatic Brain Injury](#)
- [Special Care: an Oral Health Professionals Guide to Serving Young Children with Special Health Care Needs](#)
- [Bright Futures Oral Health Pocket Guide](#)
- [American Academy of Pediatrics Oral Health Initiative](#)
- [American Academy of Pediatric Dentistry: 2008-09 Definitions, Oral Health Policies and Clinical Guidelines](#)
- [MCH Resource Center](#)
- [ASTDD-Special Needs](#)
- [Block Oral Disease, MA](#)
- [NOHIC-NIDCR publications](#)





Dear Colleague,

The Washington State Department of Health (DOH) provides print-ready files (PDFs) of health education materials. To ensure that the original quality of the piece is maintained, please read and follow the instructions below and the specifications included for professional printing.

- **Use the latest version.** DOH materials are developed using the most current information available, are checked for clinical accuracy, and are field tested with the intended audience to ensure they are clear and readable. DOH programs make periodic revisions to educational materials, so please check this web site to be sure you have the latest version. DOH assumes no responsibility for the use of this material or for any errors or omissions.
- **Do not alter.** We are providing this artwork with the understanding that it will be printed without alterations and copies will be free to the public. Do not edit the text or use illustrations or photographs for other purposes without first contacting us. Please do not alter or remove the DOH logo, publication number or revision date. If you want to use a part of this publication for other purposes, contact the Office of Health Promotion first.
- **For quality reproduction:** Low resolution PDF files are intended for black and white or color desktop printers. They work best if you are making only one or two copies. High resolution PDF files are intended for reproducing large quantities and are set up for use by professional offset print shops. The high resolution files also include detailed printing specifications. Please match them as closely as possible and insist on the best possible quality for all reproductions.

If you have questions, contact:
Office of Health Promotion
P.O. Box 47833 Olympia, WA 98504-7833
(360) 236-3736

Sincerely,
Health Education Resource Exchange Web Team